Sarcouncil Journal of Internal Medicine and Public Health

ISSN(Online): 2945-3674

Volume- 04| Issue- 06| 2025

Research Article **Received:** 01-10-2025| **Accepted:** 25-10-2025 | **Published:** 09-11-2025

Measuring Success in Physical Therapy: Beyond Pain Scales and Gait Scores

Priti Nathani

University of Montana, Missoula, MT

Abstract: Pain scales, gait scores, and other traditional measures in physical therapy are long overdue in quantifying improvements in patients. Yet, these tests only give a partial and, at best, incomplete picture of recovery, because they are specific about symptoms and motor recovery. The following review examines the more widespread, comprehensive measures of success in the physical therapy field, going beyond standard instruments. It talks of the limits of pain and gait measures, the increasing use of patient-reported outcome measures (PROMs), functional evaluation, wearables, psychosocial biomarkers, and treatment adherence. Stress is put on the incorporation of the emotional, cognitive, and social aspects into the assessment schemes, which are in line with a biopsychosocial model of care. The article suggests that a multidimensional evaluation of the results of the treatment is in favor, which takes into account not only that the symptoms have been removed but also that actual functional improvement and patient success are achieved. By referencing articles with high impact, this review puts forward an original and updated concept of quantifying the success of physical therapy.

Keywords: Physical therapy, pain scales, gait scores, patient-reported outcomes, wearable technology, functional assessment, psychosocial factors, rehabilitation success, biopsychosocial model, and treatment adherence.

INTRODUCTION

PT is central to rehabilitation, management of chronic diseases, injury, or restoration of physical functioning, or improvement of physical functions of all ages. Evaluations based on pain scales and gait scores are the traditional methods of determining the result of physical therapy. Such tools as the Visual Analogue Scale (VAS), the Numeric Pain Rating Scale (NPRS), or the measures of gait parameters (e.g., speed, cadence, the length of the stride) are popular as they are simple and easily accessible. Nevertheless, in the context in which healthcare is becoming more patient-centred and patient-driven, and hospital evaluations are more holistic, these measures are not sufficient to reflect the multidimensional aspects of recovery. Regardless of the measurable level, pain scores remain subjective and depend on such aspects as mood, culture, and pain handling approaches (Turk, D. C., & Melzack, R. 2011) Likewise, the gait scores mainly evaluate motor ability without taking into consideration contextual execution, stamina, or integration into everyday activities. Recently, an increasingly recognized literature indicates that effective rehabilitation should be able to address psychological well-being, functional independence, patient satisfaction, and quality of life (Mathis, R. A. *et al.*, 2019). New paradigms of measurement and evaluation systems are inclined toward dynamic, multi-dimensional instruments which encompass patient-reported outcomes (PROs), real-world functional scales, biological markers, psychosocial, and adherence abnormalities. It will not be satisfied with symptom relief but with the recovery of meaningful participation in roles in life. In addition to that, increased technological capabilities, including the use of sensors and artificial intelligence, have further widened the physical therapist's tool set that enables constant and personalized monitoring of the healing process. This paper gives an overview of the dynamic nature of how success in physical therapy has been measured, discussing the caveats when it comes to traditional instruments and showing some of the newer approaches, which give a more complex and patient-oriented view of the effectiveness of treatment.

LIMITATIONS OF TRADITIONAL METRICS: PAIN SCALES AND GAIT SCORES

Most frequent outcome measures used in physical therapy pain scales and gait evaluations are of limited use in gaining information about functional and psychosocial recovery. Pain as a significant problem varies day-to-day and is not necessarily related to objective functional loss or quality of life (Cleeland, C. S., & K. M. Ryan. 1994). As an example, even when the functional ability and mobility are enhanced in chronic cases of musculoskeletal disorders, their reports are of pain throughout, implying disparity between the perceived pain ratings and the practical achievements (Archer, K. R. *et al.*, 2012). Equally, gait scores like the Timed Up and Go (TUG) test, 10-Meter Walk Test, and Berg Balance Scale evaluate mainly locomotor abilities by failing to gauge fully the capacity of the individual to execute activities of daily living (ADLs), social participation, and self-efficacy. Clinical

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

***Corresponding Author:** Priti Nathani

DOI- https://doi.org/10.5281/zenodo.17566017

performance of automatically walking patterns, as well, could be different in comparison with real life because of artificial conditions or motivation (Nilsagård, Y. *et al.,* 2009). Further, this dependence on these parameters does not curb the cognitive and emotional aspects of rehabilitation. Depression, anxiety, and kinesiophobia can significantly slow progress, but this fact is not seen in ordinary scales (Vlaeyen, J. W., & S. J. Linton. 2000). It is impossible to neglect these dimensions and have a still effective treatment plan on paper, and the patient fails to perform the most common actions. The interventions are limited in tailoring by the skin-deep approach toward pain and gait. Two patients who have the same pain scores, in other words, can face extremely different functional barriers or psychosocial issues, and this may require tailored solutions. Thus, a more detailed framework is needed to recognize the effect of physical therapy and optimize it.




Figure 1: Limitations of Pain and Gait Measures in Physical Therapy

This figure highlights key limitations of relying solely on pain scales and gait assessments, including limited insight into overall recovery, a lack of evaluation of psychosocial and cognitive factors, and challenges in tailoring interventions for individual needs.

PATIENT-REPORTED OUTCOME MEASURES (PROMS)

Patient-reported outcome measures (PROMs) have come up as critical measures in determining the efficiency of physical therapy regardless of the metrics assessed. PROMs describe the patient's opinion of his or her health, function, and quality of life, which makes it possible to comprehend the results of the therapeutic work more comprehensively (Beaton, D. E. *et al.,* 2001). Among the tools that have been tested on different groups of people with different disabilities, there an instruments like the Disabilities of the Arm, Shoulder and Hand (DASH) questionnaire, the Oswestry disability Index (ODI), and the Knee Injury and Osteoarthritis outcome Score (KOOS) (Roos, E. M. *et al.,* 1998). In contrast to clinician-administered instruments, PROMs include a patient voice in an assessment, thereby corresponding to the idea of person-centered care. These instruments enable monitoring of the progress over time, recognition of outstanding needs, and determining the effects of the interventions in the day-to-day life of the patient. PROMs also play an important role in shared decision-making, which enhances collaboration between the therapist and the patient (Black, N. 2013). It has been evidenced that the use of PROMs during regular care enhances communication, boosts patient satisfaction, and treatment adherence (Valderas, J. *et al.,* 2008). Moreover, the long-term outcomes are predicted by means of PROMs as compared to certain

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher

customary indicators. What is typical is the use of the baseline SF-36 or EQ-5D scores that are usually linked to functional recovery and returning to work after an intervention (Garratt, A. M. *et al.*, 1993). Nonetheless, the administration of PROMs should be efficient and aware of language, literacy, and cultural discrepancies in order to be valid. The digitalization of the process of collecting and interpreting PROMs, as well as automated systems, is actively being used, which is why it can be deemed a convenient aspect of one of the latest and most up-to-date clinical practices.

OBJECTIVE FUNCTIONAL ASSESSMENTS AND WEARABLE TECHNOLOGY

Functional tests of real-time activities demonstrate the picture of recovery more clearly as compared to isolated clinical tests. Examples of such tools are the Functional Independence Measure (FIM), 6-Minute Walk Test (6MWT), and Short Physical Performance Battery (SPPB), which consider endurance, strength, balance, and performance in specific tasks (Studenski, S. 2011). Assessments such as these can especially be helpful in geriatrics and neurological rehabilitation, where becoming independent in the ADLs is one of the important objectives. Wearable technology has also revolutionized in functional assessment. Inertial measurement units (IMUs), accelerometers, and gyroscopes are devices that enable constant tracking of movement patterns, activity, and energy going out of the clinic (Salaorni, F. *et al.*, 2024). Such data can determine minor gains or slacks that cannot be assessed by routine visits. Wearable devices allow for providing real-time feedback, which is helpful to introduce adaptive measures and promote motivation. As an illustration, exercise count data and exercise duration would assist with personalization of an aerobic program by patients with cardiovascular diseases. Tele-rehabilitation platforms have also been developed, which combine wearable data to enable remote supervision of patients, with therapy being available in less-served areas (Cottrell, M. A., & T. G. Russell. 2020). There is also evidence in research that warns of the predictive ability of wearable-generated metrics. Step counts, cadence, and postural transitions have been associated with measures of fall risk, frailty, and recovery course in the sternal surgical (after operation) and geriatric populations (Okubo, Y. *et al.*, 2016). Nevertheless, the issues of data security, compliance, and their interpretation are the problems that should be addressed with caution to be successfully introduced into care pathways.




Figure 2: Functional Assessments and Wearable Technology in Rehabilitation

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher

This figure shows how traditional assessments (FIM, 6MWT, SPPB) and wearable technology (accelerometers, IMUs, gyroscopes) are used together to monitor endurance, strength, and balance, providing real-time feedback and predictive insights during physical therapy.

PSYCHOLOGICAL AND SOCIAL INDICATORS OF RECOVERY

The success of rehabilitation cannot be measured purely within the physical scope of recovery. Physical therapy regards emotional and psychological health as the pillars of a positive long-term outcome. Depression, anxiety, coping mechanisms, motivation, and social support are other factors that play a significant role in the relationship with therapy and the results of potential long-term effects. Unluckily enough, even conservative physical therapy measurements rarely embrace such psychosocial factors, even though a body of research is accumulating to demonstrate their relevance (Baird, C. L., & L. Sands. 2004). In recovering patients with injury or chronic illness, the psychological consequences are immense, such as fear-avoidant behaviour, frustration, or loss of identity due to reduced physical ability. The latter emotional states may increase the perception of pain, participation in treatment, and eventual functional improvement. The psychological constructs are the area where measurement tools can be applied; the Tampa Scale for Kinesiophobia (TSK), Pain Catastrophizing Scale (PCS), and Hospital Anxiety and Depression Scale (HADS) are validated devices that measure these psychological constructs and might be used together with physical tests (Sullivan, M. J. *et al.*, 1994). Another essential predictor of recovery is the social determinants of health, which entail social support, socioeconomic status, occupation, and community participation. Research has indicated that those patients who have good social support and living status exhibit improved compliance with the treatment regimen and quicker functional gains (Sveinsdottir, V. *et al.*, 2012). On the other hand, isolation, poverty, or inaccessibility to transportation can hinder engagement and distort conventional measures of recovery. Also, the collaboration of patient and therapist, known as the therapeutic alliance, is a central factor in motivation, engagement, and felt success. Good rapport, denoted by understanding, communication, and sharing of objectives, can raise morale and enhance compliance with given exercises and exercise routines (Hall, A. M. *et al.*, 2010). Direct representation of this alliance is patient satisfaction, which is gaining prominence due to its emergence as one of the most significant indicators of quality care and success in rehabilitation efforts. In order to receive a complete insight into the psychological and social aspects of recovery, clinicians are welcome to implement a biopsychosocial model. This practice not only draws biological conditions together with psychological conditions and social situations but also allows the development of personal approaches in treatment. Such a holistic assessment is facilitated by the International Classification of Functioning, Disability and Health (ICF) model, organized by the World Health Organization in terms of the interaction between health conditions and contextual factors (Cieza, A., & G. Stucki. 2008). Interdisciplinary cooperation is also needed when it comes to the treatment of psychological and social variables. Physical therapists might require signing off on psychology therapists, social workers, or case managers in complex cases where they are required to provide comprehensive care to the patient. Integrating counseling, motivational interviewing, or behavioral interventions as a part of the therapy sessions can help greatly to improve the results of work with patients who have to deal with psychosocial issues. Finally, their psychological resilience and social support not only impact the short-term objectives in rehabilitation, it also predict their long-term independence and well-being. A change towards the inclusion of such indicators into everyday working is a crucial change due to the need to assess the real triumph of physical therapy interventions.

To operationalize these psychological and contextual components within clinical settings, the following table outlines key psychosocial indicators, corresponding evaluation tools, and their clinical relevance in guiding patient-centered physical therapy outcomes.

Table 1: Psychosocial and Contextual Indicators of Physical Therapy Success

Indicator Category	Example Factors	Standardized Tools for Assessment	Clinical Implications
Cognitive-	Pain beliefs, fear	Tampa Scale for	Tailoring of behavioral

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher

Behavioral Variables	avoidance, and coping style	Kinesiophobia (TSK), Pain Catastrophizing Scale (PCS)	therapy components; identifying maladaptive responses to pain
Affective and Mood Disorders	Depression, anxiety, frustration, and emotional resilience	Hospital Anxiety and Depression Scale (HADS), Beck Depression Inventory (BDI)	Psychological referral, pacing strategies, and graded exposure for functional reintegration
Motivational and Self-Efficacy	Goal ownership, confidence in performing tasks	Self-Efficacy for Exercise Scale, Goal Attainment Scaling (GAS)	Influences adherence; guides goal-setting and treatment personalization
Socioeconomic and Environmental Context	Financial stability, employment, housing, transport	Social Determinants of Health Questionnaire (SDOHQ), Employment Support Assessment (ESA)	Coordination with social services impacts accessibility and continuity of care
Social Integration and Support	Presence of caregivers, peer networks, community ties	Medical Outcomes Study Social Support Survey (MOS-SSS)	Enhances adherence and morale; buffer against relapse or drop-out
Therapist-Patient Relationship	Empathy, communication quality, and therapeutic alliance	Working Alliance Inventory (WAI), MedRisk Satisfaction Scale (MRPS)	Determines patient satisfaction and engagement; can predict treatment retention

TREATMENT ADHERENCE, SATISFACTION, AND HOLISTIC SUCCESS

Patient satisfaction and treatment adherence are pivotal aspects of therapeutic success that are underemphasized in most cases. To a great extent, better outcomes are in direct relationship with high rates of adherence to recommended regimens such as home exercise programs and lifestyle changes. Nevertheless, the conventional evaluation, like pain or range of movement, does not consider these behaviors, although they are critical to maintaining the long-term recovery (Jack, K. *et al.*, 2010). Since compliance is a multifactorial process, it depends on the clarity of instructions, perceived utility, the burden of therapy, and how good the relationship between the therapist and the patient is. Home-based programs have low adherence, which is normally caused by a lack of supervision, improper motivation, or limited knowledge of the exercises. Newer ways of implementation like cell phone applications, distant coaching, and wristband comments, are demonstrating some success in compliance and measuring participation continuously (Schoeppe, S. *et al.*, 2017). The other effective, subjective measure of success is patient satisfaction. Although usually taken as second to objective clinical outcomes, satisfaction is the perceived quality, accessibility, communication, and accomplishment of the goals. Such dimensions can be measured in structured ways using instruments such as the Patient Satisfaction Questionnaire (PSQ) or using the MedRisk Instrument of Measuring Patient Satisfaction with Physical Therapy (MRPS). Increased satisfaction implies improved adherence, reductions in dropout outcomes, and enhanced self-reported results (Jahan, A. M. *et al.*, 2021). Holistic success is also associated with the process of putting the patients back to their former position, be it work, sporting activities, or social duties. The rates of returns-to-function and goal attainment scales (GAS) provide individualized measures to assess progress. Specifically, the GAS enables customized goal setting and result measurement that helps build a feeling of ownership and connection between a patient and a therapist (Turner-Stokes, L. 2009). Also, the long-term follow-up of performance and quality of life is required. The WHOQOL-BREF or SF-36 is among the measures that allow assessing physical, psychological, and social realms months after the end of therapy. This kind of follow-up makes sure that such gains do not occur temporarily but instead are long-lasting in the daily life of the patient. The introduction of satisfaction, adherence, and whole-of-life outcome changes the definition of success to successful symptom reduction to restoration, and the existence of a meaningful life. A more holistic assessment paradigm not only leads to superior results but also to more ethical, empathic, and patient-directed care provision.

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher

CONCLUSION

Success, then, is a matter of measure in physical therapy-but there needs to be a paradigm change in how success is measured- away from classical measures of success like pain scores and gait measures towards more enlarged metrics of success. Although the tools present convenient baseline data, they do not reflect the complexity of human recovery. The enlistment of patient-reported outcomes, objective functional measurement, wearable technology, psychosocial predictors, and adherence and satisfaction measures contributes to a more accurate, comprehensive, systematic, and significant assessment of the therapeutic performance. This mixture and amalgamation of various measures are in line with contemporary healthcare that is patient-centered and individualized. It allows physical therapists to create interventions with greater specificity, check the progress more adequately, and make patients more involved. It is expected that in future research and clinical practice, efforts will both build on these assessment frameworks and hone them, so that they succeed not just in reaching the symptom reduction focus of physical therapy but also the goals of restored function, independence, and quality of life.

REFERENCES

1. Turk, D. C., and R. Melzack. "The measurement of pain and the assessment of people experiencing pain."
2. Mathis, R. A., J. D. Taylor, B. H. Odom, and C. Lairamore. "Reliability and validity of the patient-specific functional scale in community-dwelling older adults." *Journal of Geriatric Physical Therapy* 42.3 (2019): E67–E72.
3. Cleeland, C. S., and K. M. Ryan. "Pain assessment: global use of the Brief Pain Inventory." *Annals of the Academy of Medicine, Singapore* 23.2 (1994): 129–138.
4. Archer, K. R., R. C. Castillo, S. T. Wegener, C. M. Abraham, and W. T. Obremskey. "Pain and satisfaction in hospitalized trauma patients: the importance of self-efficacy and psychological distress." *Journal of Trauma and Acute Care Surgery* 72.4 (2012): 1068–1077.
5. Nilsagård, Y., C. Lundholm, E. Denison, and L. G. Gunnarsson. "Predicting accidental falls in people with multiple sclerosis—a longitudinal study." *Clinical Rehabilitation* 23.3 (2009): 259–269.
6. Vlaeyen, J. W., and S. J. Linton. "Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art." *Pain* 85.3 (2000): 317–332.
7. Beaton, D. E., J. N. Katz, A. H. Fossel, J. G. Wright, V. Tarasuk, and C. Bombardier. "Measuring the whole or the parts? Validity, reliability, and responsiveness of the Disabilities of the Arm, Shoulder and Hand outcome measure in different regions of the upper extremity." *Journal of Hand Therapy* 14.2 (2001): 128–142.
8. Roos, E. M., H. P. Roos, L. S. Lohmander, C. Ekdahl, and B. D. Beynnon. "Knee Injury and Osteoarthritis Outcome Score (KOOS)—development of a self-administered outcome measure." *Journal of Orthopaedic & Sports Physical Therapy* 28.2 (1998): 88–96.
9. Black, N. "Patient-reported outcome measures could help transform healthcare." *BMJ* 346 (2013).
10. Valderas, J., A. Kotzeva, M. Espallargues, G. Guyatt, C. E. Ferrans, M. Y. Halyard, et al. "The impact of measuring patient-reported outcomes in clinical practice: a systematic review of the literature." *Quality of Life Research* 17 (2008): 179–193.
11. Garratt, A. M., D. A. Ruta, M. I. Abdalla, J. K. Buckingham, and I. T. Russell. "The SF36 health survey questionnaire: an outcome measure suitable for routine use within the NHS?" *British Medical Journal* 306.6890 (1993): 1440–1444.
12. Studenski, S., S. Perera, K. Patel, C. Rosano, K. Faulkner, M. Inzitari, et al. "Gait speed and survival in older adults." *JAMA* 305.1 (2011): 50–58.
13. Salaorni, F., G. Bonardi, F. Schena, M. Tinazzi, and M. Gandolfi. "Wearable devices for gait and posture monitoring via telemedicine in people with movement disorders and multiple sclerosis: a systematic review." *Expert Review of Medical Devices* 21.1–2 (2024): 121–140.
14. Cottrell, M. A., and T. G. Russell. "Telehealth for musculoskeletal physiotherapy." *Musculoskeletal Science and Practice* 48 (2020): 102193.
15. Okubo, Y., Y. Osuka, S. Jung, F. Rafael, T. Tsujimoto, T. Aiba, et al. "Walking can be more effective than balance training in fall prevention among community-dwelling older adults." *Geriatrics & Gerontology International* 16.1 (2016): 118–125.

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher

16. Baird, C. L., and L. Sands. "A pilot study of the effectiveness of guided imagery with progressive muscle relaxation to reduce chronic pain and mobility difficulties of osteoarthritis." *Pain Management Nursing* 5.3 (2004): 97–104.
17. Sullivan, M. J., S. R. Bishop, and J. Pivik. "The pain catastrophizing scale: development and validation." *Psychological Assessment* 7.4 (1995): 524.
18. Sveinsdottir, V., H. R. Eriksen, and S. E. Reme. "Assessing the role of cognitive behavioral therapy in the management of chronic nonspecific back pain." *Journal of Pain Research* (2012): 371–380.
19. Hall, A. M., P. H. Ferreira, C. G. Maher, J. Latimer, and M. L. Ferreira. "The influence of the therapist–patient relationship on treatment outcome in physical rehabilitation: a systematic review." *Physical Therapy* 90.8 (2010): 1099–1110.
20. Cieza, A., and G. Stucki. "The International Classification of Functioning Disability and Health: its development process and content validity." *European Journal of Physical and Rehabilitation Medicine* 44.3 (2008): 303–313.
21. Jack, K., S. M. McLean, J. K. Moffett, and E. Gardiner. "Barriers to treatment adherence in physiotherapy outpatient clinics: a systematic review." *Manual Therapy* 15.3 (2010): 220–228.
22. Schoeppe, S., S. Alley, A. L. Rebar, M. Hayman, N. A. Bray, W. Van Lippevelde, et al. "Apps to improve diet, physical activity, and sedentary behaviour in children and adolescents: a review of quality, features, and behaviour change techniques." *International Journal of Behavioral Nutrition and Physical Activity* 14 (2017): 1–10.
23. Jahan, A. M., A. E. Rwaiha, S. R. Gusaibat, N. A. Al-Ahwal, Z. M. Al-Jafairi, and M. A. Al-Rashidi. "Patient satisfaction with physiotherapy services in Libya: a cross-sectional study." *Journal of Patient Experience* 8 (2021): 23743735211033195.
24. Turner-Stokes, L. "Goal attainment scaling (GAS) in rehabilitation: a practical guide." *Clinical Rehabilitation* 23.4 (2009): 362–370.

Source of support: Nil; **Conflict of interest:** Nil.

Cite this article as:
Nathani, P. "Measuring Success in Physical Therapy: Beyond Pain Scales and Gait Scores." *Sarcouncil Journal of Internal Medicine and Public Health* 4.6 (2025): pp 16-22.

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher